

## ACKNOWLEDGEMENTS

We should like to thank our seafaring colleagues Derek Smith and Rod Grout for providing extra material. Our mentor throughout this project, R. B. Sibson of Auckland, is sincerely thanked for his continuing assistance with our seabird recording.

## LITERATURE CITED

- BARTLE, J. A. 1974. Seabirds of eastern Cook Strait, New Zealand, in autumn. *Notornis* 21: 135-166.  
 FOWLER, J. A. 1973. A transect diagram for oceanic birds recorded between McMurdo Sound, Antarctica, and New Zealand, February-March 1973. *Notornis* 20: 314-317.  
 SUMMERHAYES, C. P. 1967. Seabirds seen in the northern Tasman Sea in winter. *N.Z. Jour. Freshwater Res.* 3: 560-570.

NEIL CHESHIRE, 140 Glamorgan Drive, Torbay, Auckland; JOHN JENKINS, 14 Lochiel Road, Remuera, Auckland; PETER NESFIELD, 8 Opuia Street, Belmont, Auckland.



## SHORT NOTES

## WHITE-FACED HERONS ON CHATHAM ISLAND

On 9 January 1978, while in the south-west of Chatham Island with David Crockett's successful 'Taiko' Expedition, my husband and I were traversing the coast from just north of the Little Awatotara Creek to north of the Durham Point area. While high on a bluff, we saw a White-faced Heron (*Ardea novaehollandiae*) fly out from a ledge below us, and we saw a nest.

The nest, some 10 m above sea-level, was in a small cleft. It contained a few pieces of driftwood and seaweed, and two sky-blue eggs. The rock face below was white with droppings.

On 21 January we revisited the site and flushed the bird — two eggs; and twice on the 29th — one chick and one egg. As we then had to leave the island no further observations were possible.

Could it be that this site was chosen because of the complete lack of trees more than 3 m high within a 1500 m radius?

AUDREY GORDON, 4 Lewis Street, Kamo, Northland.



## AN INTERESTING BAND RECOVERY

On 21 April 1978, I found a juvenile New Zealand Black-browed Mollymawk (*Diomedea melanophris impavida*) carrying a New Zealand band No. 22519, about 3 km north of Himitangi, on the Wellington west coast. The bird, which had probably died within 24 hours, had been banded by the Campbell Island meteorological party only seven days earlier, in its nest on Bull Rock, Campbell Island — 1470 km away.

HUGH A. ROBERTSON, No. 1 R.D., Aokautere Drive, Palmerston Nth.