

SHORT NOTE

An unusual last meal for a fairy prion (titi wainui, *Pachyptila turtur*)

MICHAEL WAKELIN
47 Hunt Street, Dunedin, New Zealand

On the 28 February 2023 a dead fairy prion (titi wainui, *Pachyptila turtur*) was found washed up on Saint Clair beach, Ōtepoti Dunedin (45.909580°S, 170.502050°E). The bird appeared to have been dead for at least one week and was placed in a covered sand box to continue its decay. When inspected six months later the abdominal cavity was found to contain the elytra of 22 eucalyptus tortoise beetles (*Paropsis charybdis* Stål, 1860).

These chrysomelid beetles are approximately 10 mm long, hemispherical, and a mottled tan/brown/pink colour. They were accidentally introduced to New Zealand in the 1920's, becoming widespread over 50 years ago. As a herbivorous pest of *Eucalyptus* gum trees it has been the focus of biocontrol research (White 1973; Radics *et al.* 2018). The adult beetles are strong and widespread fliers in late summer (McGregor 1989; Selman 1994) and many were observed in late February 2023 at the Hump Ridge, Southland (R. Goldsmith *pers. comm.*).

Flying beetles could be blown offshore or swept down rivers and may concentrate along convergent zones of coastal currents. Previously a large number were found washed up on Bay of Plenty (North Island, New Zealand) beaches (White 1973).

Gum trees in coastal Otago are relatively common with over 12,000 hectares of plantation *Eucalyptus* in Otago/Southland in 2014 (SWC 2015). Fairy prions are also relatively common in coastal Otago and breed on the Otago coast including at Saint Clair and Green Island.

Fairy prions usually feed on euphausiids, amphipods, cephalopods, fish, and molluscs but will readily feed on anything resembling food on the water surface (Harper & Fowler 1987), particularly if the bird is starved or a juvenile (Harper & Fowler 1987; Acampora *et al.* 2014). The bird in this observation was a fledgling as indicated by the incomplete fusion of some of the bones (N. Rawlence *pers. comm.*). Terrestrial insects have occasionally been reported in seabird stomach contents (Steele & Klages 1986; Gartshore *et al.* 1988; Petry *et al.* 2008; Acampora *et al.* 2014) but may be considered an irregular and unimportant dietary

component (Gartshore *et al.* 1988) and therefore omitted from dietary study results.

Although low numbers of the eucalyptus tortoise beetles have been recorded in the stomachs of rooks (*Corvus frugilegus*) (Porter 1979), common starlings (*Sturnus vulgaris*) (Moeed 1980), house sparrows (*Passer domesticus*) and European greenfinches (*Chloris chloris*) (MacMillan 1981), the beetle larvae and adults are considered to be poisonous or at least unpalatable (Moore 1967; Selman 1994). It is not known if eating 22 *Paropsis charybdis* beetles caused the death of this prion.

ACKNOWLEDGEMENTS

Support, suggestions, information, and improvements were provided by Bruce McKinlay, Graeme Loh, Nic Rawlence, Ruth Goldsmith, Karen Reader, James Tweed, and an anonymous reviewer.

LITERATURE CITED

- Acampora, H.; Schuyler, Q.; Townsend, K.A.; Hardesty, B.D. 2014. Comparing plastic ingestion in juvenile and adult stranded short-tailed shearwaters (*Puffinus tenuirostris*) in eastern Australia. *Marine Pollution Bulletin* 78: 63–68.
- Gartshore, N.A.; Steele, W.K.; Klages, N.T. 1988. Summer diet of the Salvin's prion at sub-Antarctic Marion Island. *South African Journal of Zoology* 23: 309–313.
- Harper, P.C.; Fowler, J.A. 1987. Plastic pellets in New Zealand storm-killed prions (*Pachyptila* spp.) 1958–1977. *Notornis* 34: 65–70.
- MacMillan, B.W.H. 1981. Food of house sparrows and greenfinches in a mixed farming district, Hawke's Bay, New Zealand. *New Zealand Journal of Zoology* 8: 93–104.
- McGregor, P.G. 1989. Ecology of *Paropsis charybdis* Stal (Coleoptera: Chrysomelidae): A Eucalyptus defoliator in New Zealand. Unpubl. PhD

thesis, Massey University, Palmerston North, New Zealand.

- Moeed, A. 1980. Diets of adult and nestling starlings (*Sturnus vulgaris*) in Hawke's Bay, New Zealand. *New Zealand Journal of Zoology* 7: 247–256.
- Moore, B. 1967. Hydrogen cyanide in the defensive secretions of larval Paropsini (Coleoptera: Chrysomelidae). *Australian Journal of Entomology* 6: 36–38.
- Petry, M.V.; da Silva Fonseca, V.S.; Krüger-Garcia, L.; da Cruz Piuco, R.; Brummelhaus, J. 2008. Shearwater diet during migration along the coast of Rio Grande do Sul, Brazil. *Marine Biology* 154: 613–621.
- Porter, R.E.R. 1979. Food of the rook (*Corvus frugilegus* L.) in Hawke's Bay, New Zealand. *New Zealand Journal of Zoology* 6: 329–337.
- Radics, R.I.; Withers, T.M.; Meason, D.F.; Stovold, T.; Yao, R. 2018. Economic impact of eucalyptus tortoise beetle (*Paropsis charybdis*) in New Zealand. Scion (New Zealand Forest Research Institute Ltd.), Rotorua, New Zealand. 26 pp.
- Selman, B.J. 1994. The biology of the paropsine eucalyptus beetles of Australia. Pp. 555–565. *In*: Jolivet, P.H.; Cox, M.L.; Petitpierre, E. (ed.). Novel aspects of the biology of the Chrysomelidae. Amsterdam: Kluwer Academic.
- Steele, W.K.; Klages, N.T. 1986. Diet of the blue petrel at sub-Antarctic Marion Island. *South African Journal of Zoology* 21: 253–256.
- SWC 2015. <https://www.southernwoodcouncil.co.nz/wp-content/uploads/2015/11/Otago-Southland-Forestry-Profile-2015-03.pdf>
- White, T.C.R. 1973. The establishment, spread and host range of *Paropsis charybdis* Stal (Chrysomelidae) in New Zealand. *Pacific Insects* 15: 59–66.

Keywords: Fairy prion, titi wainui, *Pacyptila turtur*, eucalyptus tortoise beetle, *Paropsis charybdis*, stomach contents